

Adelic Formulas for Gamma- and Beta-functions in Algebraic Numbers Fields

V. S. Vladimirov

(Steklov Mathematical Institute, Moscow, Russia)

Abstract. On the basis of analysis on the adèle ring of any algebraic numbers field (Tate's formula) a regularization for divergent adelic products of gamma- and beta-functions for all completions of this field are proposed, and corresponding regularized adelic formulas are obtained.

§1. Introduction

Freund and Witten [1](1987) and, independently a little later, Volovich [2] (1988), suggested for the beta-function B_∞ of the real numbers field $\mathbb{R}$ the following adelic formula

$$B_\infty(\alpha, \beta) \prod_{p=2}^{\infty} B_p(\alpha, \beta) = 1. \quad (1.1)$$

where B_p is the beta-function of p-adic numbers field $\mathbb{Q}_p$.

However, the infinite product in (1.1) diverges for all α and β and, therefore, it is not clear how to understand Eq. (1.1). A detailed discussions of these matters can be found in [3]–[8].

In this paper on the basis of analysis on the adèle ring (Tate's formula) for any algebraic numbers field a regularization for divergent infinite products of gamma- and beta-functions for all characters (ramified or non) is proposed, and the regularized adelic formulas are derived.

More precisely, let $\mathbb{K} = \mathbb{Q}(\epsilon)$ be an algebraic numbers field of degree $n = \sigma + 2\tau$ where σ and 2τ are numbers of real and complex roots resp. of the minimal polynomial for a primitive element ϵ of field $\mathbb{K}$. Then the

following regularized adelic formulas for gamma- and beta-functions are valid (see Eqs (4.18) and (5.4))

$$\prod_{v=1}^{\sigma} \Gamma_{\infty}(\alpha + i\alpha_v; \nu_v) \prod_{v=\sigma+1}^{\sigma+\tau} \Gamma_{-\infty}(\alpha + i\alpha_v; \nu_v) \operatorname{reg} \prod_{v \in F} \Gamma_{q_v}(\alpha + i\alpha_v) \\ = \kappa \omega(C) [|D|N(J)]^{1/2-\alpha}, \quad (1.2)$$

$$\prod_{v=1}^{\sigma} B_{\infty}(\alpha + i\alpha_v, \nu_v; \beta + i\beta_v, \mu_v) \prod_{v=\sigma+1}^{\sigma+\tau} B_{-\infty}(\alpha + i\alpha_v, \nu_v; \beta + i\beta_v, \mu_v) \\ \times \operatorname{reg} \prod_{v \in F} B_{q_v}(\alpha + i\alpha_v, \beta + i\beta_v) = \kappa \sqrt{|D|N(J)}. \quad (1.3)$$

where $\Gamma_{\infty}(\alpha; \nu)$ and $B_{\infty}(\alpha, \nu; \beta, \mu)$ are gamma- and beta-functions of the real numbers field $\mathbb{Q}_{\infty} = \mathbb{R}$ for the characters (see §2)

$$\theta_v(x) = \operatorname{sgn}^{\nu} x, \quad \theta'_v(x) = \operatorname{sgn}^{\mu} x,$$

$$\nu = \nu_v, \mu = \mu_v \in F_2, v = 1, 2, \dots, \sigma;$$

$B_{-\infty}(\alpha, \nu; \beta, \mu)$ is the beta-function of the complex numbers field $\mathbb{Q}_{-\infty} = \mathbb{C}$ for characters (see §2)

$$\theta_v(x) = x^{\nu} (x\bar{x})^{-\nu/2}, \quad \theta'_v(x) = x^{\mu} (x\bar{x})^{-\mu/2},$$

$$\nu = \nu_v, \mu = \mu_v \in Z, \quad v = \sigma + 1, \dots, \sigma + \tau;$$

and

$$B_{q_v}(\alpha, \beta) = \Gamma_{q_v}(\alpha) \Gamma_{q_v}(\beta) \Gamma_{q_v}(\gamma), \quad \alpha + \beta + \gamma = 1 \quad (1.4)$$

is the beta-function of a local p-field $\mathbb{K}_v$ with module q_v for unramified characters, $v \in F$ (i.e. $v = \sigma + \tau + 1, \dots, \theta_v(x) \equiv 1$). Here

$$\Gamma_{q_v}(\alpha) = \frac{1 - q_v^{\alpha-1}}{1 - q_v^{-\alpha}}, \quad \Gamma_{q_v}(\alpha) \Gamma_{q_v}(1 - \alpha) = 1 \quad (1.5)$$

is the reduced gamma-function of the field $\mathbb{K}_v$ (see §2).

For further notations in (1.2) and (1.3) see §4 and §5.

For unramified characters formulas (4.19) and (5.5) (special cases of (1.2) and (1.3)) were obtained in [9],[10]. Their concrete realizations have been made in the following cases: for rational numbers field $\mathbb{K} = \mathbb{Q}$ – in [11], for quadratic fields $\mathbb{K} = \mathbb{Q}(\sqrt{d})$ – in [12], for m -circular fields $\mathbb{K} = \mathbb{Q}(\exp(2\pi i/m))$ – in [10],[9], for some cubic fields $\mathbb{K} = \mathbb{Q}(\sqrt[3]{d})$ – in [10].

Adelic formula (1.3) is applied to the four-point Veneziano and Virasoro-Shapiro amplitudes and their generalizations which connect them with string (open or closed resp.) amplitudes (see [1]–[12]).

In this paper we use notations of [10]. In particular,

$$\delta(|x| - q^\gamma) = \begin{cases} 1, & |x| = q^\gamma \\ 0, & |x| \neq q^\gamma \end{cases}, \quad \Omega(q^{-\gamma}|x|) = \begin{cases} 1, & |x| \leq q^\gamma \\ 0, & |x| > q^\gamma \end{cases}.$$

Z is the ring of rational integers; Z_+ is the set of natural numbers.

Necessary informations on number theory and analysis can be found in books [13]–[20].

The results of this paper were published shortly in Doklady RAN [21].

This work is supported in part by RFFI Grant 96-01-01008.

§2. Local fields of zero characteristic

Such fields are well-known: they are $\mathbb{R}, \mathbb{C}$ and algebraic extensions of $\mathbb{Q}_p$ (p -fields [13]). We denote them by $\mathbb{K}$.

Let x be an element of $\mathbb{K}$, $|x|$ be a valuation on $\mathbb{K}$, dx be the normed Haar measure on $\mathbb{K}$. For p -field $\mathbb{K}$ let q be its module, $\mathbb{Z}$ be the ring of integer elements, $\mathbb{Z}^\times$ be the multiplicative subgroup of $\mathbb{Z}$, I be its maximal ideal.

Let $\chi(x)$ be an additive character of $\mathbb{K}$, r be its *rank* (order) [13],[18]. (For p -field $\mathbb{K}$ the rank r is max of $\gamma \in Z$ for which $\chi(x) \equiv 1, |x| \leq q^\gamma$.) Therefore the measure $q^{-r/2}dx$ is self-dual with respect to this character.

Let

$$\omega(x) = \theta(x)|x|^{i\alpha}, \quad \alpha \in \mathbb{R} \quad (2.1)$$

be a multiplicative character of $\mathbb{K}$, ρ be its *rank* [13],[18]. (For p -field $\mathbb{K}$ the rank ρ is min of $\gamma \in Z$ for which $\omega(x) \equiv 1, x \in \mathbb{Z}, 1 - x \in I^\gamma$).

Now we introduce the following definitions [18].

Gamma-function $\Gamma(\alpha; \theta)$ of field $\mathbb{K}$ for a character θ is the analytic continuation from domain $\text{Re } \alpha > 0$ of integral

$$\Gamma(\alpha; \theta) = q^{-r/2} \int_{\mathbb{K}} \theta(x)|x|^{\alpha-1} \chi(x) dx. \quad (2.2)$$

Beta-function $B(\alpha, \theta; \beta, \theta')$ of field $\mathbb{K}$ for characters θ and θ' is the analytic continuation from tube domain $\text{Re } \alpha > 0, \text{Re } \beta > 0, \text{Re}(\alpha + \beta) < 1$ of integral

$$B(\alpha, \theta; \beta, \theta') = \theta(-1)\theta'(-1) \int_{\mathbb{K}} \theta(x)|x|^{\alpha-1} \theta'(1-x)|1-x|^{\beta-1} dx. \quad (2.3)$$

The Mellin transform $\Phi(\alpha; \theta)$ of a function $\varphi \in \mathcal{S}(\mathbb{K})$ with respect to a character θ of field $\mathbb{K}$ is the analytic continuation from domain $\text{Re } \alpha > 0$ of integral

$$\Phi(\alpha; \theta) = \int_{\mathbb{K}^\times} \varphi(x)\theta(x)|x|^\alpha d^\times x \quad (2.4)$$

where $d^\times x$ is the normed Haar measure on $\mathbb{K}^\times$.

The following formulas are valid:

$$B(\alpha, \omega; \beta, \omega') = B(\alpha + i\alpha', \theta; \beta + i\beta', \theta')$$

if $\omega'(x) = \theta'(x)|x|^{i\alpha'}$;

$$B(\alpha, \theta; \beta, \theta') = q^{r/2} \Gamma(\alpha; \theta) \Gamma(\beta; \theta') \Gamma(\gamma; \theta'') \quad (2.5)$$

if $\alpha + \beta + \gamma = 1, \theta\theta'\theta'' = 1$;

$$\Gamma(\alpha; \theta) \Gamma(1 - \alpha; \bar{\theta}) = \theta(-1), \quad \Gamma(\alpha; \omega) = \Gamma(\alpha + i\alpha'; \theta); \quad (2.6)$$

$$F[\theta(x)|x|^{\alpha-1}](\xi) = \Gamma(\alpha; \theta)\bar{\theta}(\xi)|\xi|^{-\alpha} \quad (2.7)$$

where $F[f] = \tilde{f}$ is the Fourier transform of $f \in \mathcal{S}(\mathbb{K})$ with respect to the character $\chi(x)$ and the self-dual measure $q^{-r/2}dx$;

$$\Phi(\alpha; \theta) = \Gamma(\alpha; \theta)\tilde{\Phi}(1 - \alpha; \bar{\theta})\theta(-1), \quad \Phi(\alpha; \omega) = \Phi(\alpha + i\alpha'; \theta) \quad (2.8)$$

where $\tilde{\Phi}$ is the Mellin transform of $\tilde{\varphi}$ with respect to the character $\bar{\theta}$.

In special local fields the quantities just introduced have the following particular form.

Field $\mathbb{K} = \mathbb{Q}_\infty = \mathbb{R}$. Here dx is the Lebesgue measure, $d^\times x = |x|^{-1}dx$, $q = 1$, $\chi(x) = \exp(-2\pi ix)$, $r = 0$,

$$\theta_\nu(x) = \text{sgn}^\nu x, \quad \nu \in F_2, \quad \rho = 0.$$

(F_2 means the field of residues of mod 2, i.e. it consists of elements 0 and 1.)

$$\Gamma_\infty(\alpha; \theta_\nu) = \Gamma_\infty(\alpha; \nu), \quad \nu \in F_2, \quad (2.9)$$

where

$$\Gamma_\infty(\alpha; \nu) = i^{-\nu} \pi^{1/2-\alpha} \frac{\Gamma(\frac{\alpha+\nu}{2})}{\Gamma(\frac{1-\alpha+\nu}{2})}, \quad \Gamma_\infty(\alpha; \nu)\Gamma_\infty(1 - \alpha; \nu) = (-1)^\nu; \quad (2.10)$$

and

$$B_\infty(\alpha, \theta_\nu; \beta, \theta'_\mu) = B_\infty(\alpha, \nu; \beta, \mu), \quad \nu, \mu \in F_2, \quad (2.11)$$

where

$$B_\infty(\alpha, \nu; \beta, \mu) = \Gamma_\infty(\alpha; \nu)\Gamma_\infty(\beta; \mu)\Gamma_\infty(1 - \alpha - \beta; -\nu - \mu). \quad (2.12)$$

Owing to (2.12) B_∞ is symmetric under permutations of points (α, ν) , (β, μ) , (γ, η) on the manifold

$$\alpha + \beta + \gamma = 1, (\alpha, \beta, \gamma) \in \mathbb{C}^3, \quad \nu + \mu + \eta = 0, (\nu, \mu, \eta) \in F_2^3. \quad (2.13)$$

Field $\mathbb{K} = \mathbb{Q}_{-\infty} = \mathbb{C}$. Here $|x| = x\bar{x}$, $dx = |\partial x \wedge \bar{\partial} x|$, $d^\times x = (x\bar{x})^{-1}dx$, $q = 1$, $\chi(x) = \exp(-2\pi i(x + \bar{x}))$, $r = 0$,

$$\theta_\nu(x) = x^\nu (x\bar{x})^{-\nu/2}, \quad \nu \in Z, \quad \rho = 0,$$

$$\Gamma_{-\infty}(\alpha; \theta_\nu) = \Gamma_{-\infty}(\alpha; \bar{\theta}_\nu) = \Gamma_{-\infty}(\alpha; \nu) = \Gamma_{-\infty}(\alpha; -\nu) \quad (2.14)$$

where

$$\Gamma_{-\infty}(\alpha; \nu) = i^{-|\nu|} (2\pi)^{1-2\alpha} \frac{\Gamma(\alpha + |\nu|/2)}{\Gamma(1 - \alpha + |\nu|/2)},$$

$$\Gamma_{-\infty}(\alpha; \nu) \Gamma_{-\infty}(1 - \alpha; \nu) = (-1)^\nu; \quad (2.15)$$

and

$$B_{-\infty}(\alpha, \theta_\nu; \beta, \theta_\mu) = B_{-\infty}(\alpha, \nu; \beta, \mu) \quad (2.16)$$

where

$$B_{-\infty}(\alpha, \nu; \beta, \mu) = \Gamma_{-\infty}(\alpha; \nu) \Gamma_{-\infty}(\beta; \mu) \Gamma_{-\infty}(1 - \alpha - \beta; -\nu - \mu). \quad (2.17)$$

Owing to (2.17) $B_{-\infty}$ is symmetric under permutations of points (α, ν) , (β, μ) , (γ, η) on the manifold

$$\alpha + \beta + \gamma = 1, (\alpha, \beta, \gamma) \in \mathbb{C}^3, \quad \nu + \mu + \eta = 0, (\nu, \mu, \eta) \in Z^3. \quad (2.18)$$

Fields $\mathbb{K} = \mathbb{Q}_p(\epsilon)$, $p = 2, 3, 5, \dots$. Here dx is the Haar measure on $\mathbb{K}$ normed by condition

$$\int_{|x| \leq 1} dx = 1;$$

$$d^\times x = (1 - q^{-1})^{-1} |x|^{-1} dx, \quad \int_{|x|=1} d^\times x = 1$$

where $q = p^f$, $f \in Z_+$ is the module of $\mathbb{K}$;

$$\chi(x) = \exp(2\pi i \{x\}_p)$$

where $\{x\}_p$ is the rational part of $x \in \mathbb{K}$. A multiplicative character $\theta(x)$ on $\mathbb{K}$ we normalize by the condition

$$\theta(\pi) = 1 \quad (2.19)$$

where π is the *generating element* of $\mathbb{K}$.

For unramified characters ($\theta \equiv 1, \rho = 0$) we have (cf. [10])

$$\Gamma(\alpha; 1) = q^{-r/2} \int_{|x| \leq q^{r+1}} |x|^{\alpha-1} \chi(x) dx = q^{(\alpha-1/2)r} \Gamma_q(\alpha) \quad (2.20)$$

where $\Gamma_q(\alpha)$ is the reduced gamma-function of field $\mathbb{Q}_p(\epsilon)$ with module q defined by the formula (1.5).

(The Eq. (2.20) follows directly from the p -adic integral

$$\int_{|x| \leq q^\gamma} \chi(\xi x) dx = q^\gamma \Omega(q^{\gamma-r} |\xi|) \quad (2.21).$$

Indeed, it is trivial for $|\xi| \leq q^{r-\gamma}$. Let ξ be such that $|\xi| > q^{r-\gamma}$. There exists $x_0, |x_0| = q^{r+1}$ such that $\chi(\xi x_0) \neq 0$. Then

$$\int_{|x| \leq q^\gamma} \chi(\xi x) dx = \int_{|x+x_0| \leq q^\gamma} \chi(\xi(x+x_0)) dx = \chi(\xi x_0) \int_{|x| \leq q^\gamma} \chi(\xi x) dx,$$

and (2.21) follows.)

From (2.20) it follows that $\Gamma(\alpha; 1)$ is the Mellin transform of function [10]

$$(1 - q^{-1}) q^{-r/2} \chi(x) \Omega(q^{-r-1} |x|) \in \mathcal{S}(\mathbb{K}). \quad (2.22)$$

Besides,

$$B(\alpha, 1; \beta, 1) = B_q(\alpha, \beta) = \Gamma_q(\alpha) \Gamma_q(\beta) \Gamma_q(1 - \alpha - \beta). \quad (2.23)$$

For ramified characters ($\theta \not\equiv 0, \rho \geq 1$) we have [13]

$$\Gamma(\alpha; \theta) = q^{-r/2} \int_{|x|=q^{r+\rho}} \theta(x) |x|^{\alpha-1} \chi(x) dx = \kappa(\theta) q^{(\alpha-1/2)(r+\rho)} \quad (2.24)$$

where

$$\kappa(\theta) = q^{\rho/2} \int_{|x|=1} \theta(x) \chi(\pi^{-r-\rho} x) dx, \quad |\kappa| = 1. \quad (2.25)$$

From (2.24) it follows that $\Gamma(\alpha; \theta)$ is the Mellin transform of function

$$(1 - q^{-1}) q^{-r/2} \chi(x) \delta(|x| - q^{r+\rho}) \in \mathcal{S}(\mathbb{K}). \quad (2.26)$$

§3. Analysis on the adele ring

Let $\mathbb{K} = \mathbb{Q}(\epsilon)$ be an algebraic numbers field of degree $n = \sigma + 2\tau$, and $\mathbb{Z}$ is its *ring of integer elements* (the maximal order). We denote

$$(\sigma_v, \mathbb{K}_v, \quad v = 1, 2, \dots)$$

places of field $\mathbb{K}$ [13]. Here $\mathbb{K}_v$ is $\mathbb{R}$ for infinite real places $v = 1, 2, \dots, \sigma$; $\mathbb{K}_v$ is $\mathbb{C}$ for infinite imaginary places $v = \sigma + 1, \dots, \sigma + \tau$; $\mathbb{K}_v$ is p-field $\mathbb{Q}_p(\sigma_v(\epsilon))$ for finite plces $v = \sigma + \tau + 1, \dots$

We equip quantities of §2 related to field $\mathbb{K}_v$ with index v , namely: $x_v, |x_v|_v, d_v x_v, d_v^\times x_v, \chi_v(x_v), r_v$. Let

$$\omega_v(x_v) = \theta_v(x_v) |x_v|_v^{i\alpha_v}, \quad \alpha_v \in \mathbb{R} \quad (3.1)$$

be a multiplicative character on $\mathbb{K}_v$, and ρ_v is its rank. For p-field $\mathbb{K}_v$ ($v = \sigma + \tau + 1, \dots$) let $q_v = p^{f_v}, f_v \in \mathbb{Z}_+$ be its module, $\mathbb{Z}_v$ be its ring of integer elements, $\mathbb{Z}_v^\times$ be multiplicative subgroup, I_v be maximal ideal of ring $\mathbb{Z}_v$, and $\Pi_v = \mathbb{Z} \cap I_v$ be a prime ideal of the ring $\mathbb{Z}$.

Let A be the ring of adeles,

$$X = (x_1, x_2, \dots, x_v, \dots), x_v \in \mathbb{K}_v, x_v \in \mathbb{Z}_v, v \geq V > \sigma + \tau,$$

and $A^\times$ be the group of ideles,

$$X = (x_1, x_2, \dots, x_v, \dots, x_v \in \mathbb{K}_v^\times, x_v \in \mathbb{Z}_v^\times, v \geq V > \sigma + \tau)$$

of the field $\mathbb{K}$.

Any character $\chi(X)$ of ring A is represented in the form

$$\chi(X) = \prod_v \chi_v(x_v) \quad (3.2)$$

where $\chi_v(x_v)$ is the restriction of $\chi(X)$ on subring of adeles of the form $(0, 0, \dots, x_v, 0, \dots)$.

We suppose that the character $\chi(X)$ is trivial on $\mathbb{K}$ considered as subring of A of principal adeles

$$\mathbf{x} = (\sigma_v(x), v = 1, 2, \dots, x \in \mathbb{K}),$$

i.e. $\chi(X)$ is a character of $A/\mathbb{K}$. As an example of such characters is the character

$$\chi(X) = \prod_{v=1}^{\infty} \chi_p(\mathrm{Sp}_v x_v).$$

Any character $\omega(X)$ of the group $A^\times$ is represented in the form

$$\omega(X) = \prod_v \omega_v(x_v) = \theta(X) \prod_v |x_v|_v^{i\alpha_v}, \quad \theta(X) = \prod_v \theta_v(x_v) \quad (3.3)$$

where $\omega_v(x_v)$ is defined by Eq. (3.1). It is the restriction of $\omega(X)$ on subgroup of ideles of the form $(1, 1, \dots, x_v, 1, \dots)$.

We suppose that the character $\omega(X)$ is trivial on $\mathbb{K}^\times$ of principal ideles, i.e. $\omega(X)$ is a character of $A^\times/\mathbb{K}^\times$.

There are necessary and sufficient conditions that a character $\omega(X)$ of $A^\times$ is trivial on $\mathbb{K}^\times$ [13]. For instance, for field $\mathbb{K} = \mathbb{Q}$ these conditions can be expressed in the explicit form [18]: *a character*

$$\omega(X) = \theta(X) |x|^{i\alpha} \prod_{p=2}^{\infty} |x_p|_p^{i\alpha_p}, \quad (3.4)$$

where

$$\theta(X) = \mathrm{sgn}^\nu x \prod_{p=2}^{\infty} \theta_p(x_p), \quad \theta_p(p) = 1, \quad (3.5)$$

is trivial on $\mathbb{Q}^\times$ iff

$$\theta(-1) = 1, \quad p^{i\alpha_p} = p^{i\alpha}\theta(p). \quad (3.6)$$

We recall Eqs [13],[14]

$$\prod_{v>\sigma+\tau} q_v^{r_v} = |D|, \quad N(J) = \prod_{v \in R} q_v^{\rho_v}. \quad (3.7)$$

where D is discriminant of field $\mathbb{K}$ and $N(J)$ is the norm of the *principal ideal*

$$J = \prod_{v \in R} \Pi_v^{\rho_v}$$

of the character ω ; R is the set of finite ramified places of $\mathbb{K}$. The Haar measure

$$\prod_{v=1}^{\infty} q_v^{-r_v/2} d_v x_v = |D|^{-1/2} dX \quad (3.8)$$

on A is self-dual with respect to the additive character (3.2) of ring A .

Let $V > \sigma + \tau$, and

$$\varphi(X) = \prod_{v \in F_V} \varphi_v(x_v) \prod_{v \in R} \varphi_v(x_v) \prod_{v \geq V, v \in F} \Omega(|x_v|_v), \quad \varphi_v \in \mathcal{S}(\mathbb{K}_v) \quad (3.9)$$

be a standard Schwartz-Bruchat function on A that is $\varphi \in \mathcal{S}(A)$ [13]. Here F is the set of finite unramified places of field $\mathbb{K}$, and $F_V = [v \in F : v < V]$.

The Mellin transform $\Phi(\alpha; \omega)$ of function $\varphi \in \mathcal{S}(A)$ with respect to a character ω of group $A^\times / \mathbb{K}^\times$ is the analytic continuation from domain $\text{Re } \alpha > 1$ of the integral

$$\Phi(\alpha; \omega) = \int_{A^\times} \varphi(X) \omega(X) |X|^\alpha d^\times X \quad (3.10)$$

where $d^\times X$ is the Haar measure on $A^\times$ defined by the local Haar measures $d_v^\times x_v$ of fields $\mathbb{K}_v$ (§2),

$$d^\times X = \prod_v d_v^\times x_v = \prod_{v=1}^{\sigma+\tau} |x_v|_v^{-1} d_v x_v \prod_{v>\sigma+\tau} (1 - q_v^{-1})^{-1} |x_v|_v^{-1} d_v x_v.$$

The following theorem is valid [13].

Theorem (Tate). *For unramified character ω function $\Phi(\alpha; \omega)$ is holomorphic on α everywhere except simple poles $\alpha = 0$ and $\alpha = 1$ with residues $C\Phi(0)$ and $C'\tilde{\Phi}(0)$ resp. where numbers C and C' depend only on field $\mathbb{K}$. For ramified character ω function $\Phi(\alpha; \omega)$ is entire. In addition the following relation is valid,*

$$\Phi(\alpha; \omega) = \tilde{\Phi}(1 - \alpha; \bar{\omega}) \quad (3.11)$$

where $\tilde{\Phi}$ is the Mellin transform of the Fourier transform $\tilde{\varphi}(\Xi)$ of function $\varphi(X)$ with respect to character $\bar{\omega} = \omega^{-1}$.

In detail the Tate formula (3.11) for function φ of form (3.9) can be rewritten as follows

$$\begin{aligned} \prod_{v \in F_V} \Phi_v(\alpha; \omega_v) \prod_{v \in R} \Phi_v(\alpha; \omega_v) L_V(\alpha; \omega) = \\ \prod_{v \in F_V} \tilde{\Phi}_v(1 - \alpha; \bar{\omega}_v) \prod_{v \in R} \tilde{\Phi}_v(1 - \alpha; \bar{\omega}_v) L_V(1 - \alpha; \bar{\omega}). \end{aligned} \quad (3.12)$$

In (3.12) L_V is the V -truncated Dirichlet L -function,

$$L_V(\alpha; \omega) = \prod_{v \geq V, v \in F} [1 - \lambda(v) q_v^{-\alpha}]^{-1}, \quad \lambda(v) = q_v^{-i\alpha_v} = \omega_v(\pi_v). \quad (3.13)$$

Recall that the Dirichlet L -function $L(\alpha; \omega)$ is defined by Eq.

$$L(\alpha; \omega) = \prod_{v \in F} [1 - \lambda(v) q_v^{-\alpha}]^{-1}, \quad \text{Re} > 1. \quad (3.14)$$

§4. Regularized adelic formulas for gamma-functions

Our goals are to describe a regularization and to sketch a proof of formula (1.2) for gamma-functions. To this end, we specify the Tate formula (3.12) by a special choice of standard function $\varphi(X)$ (see (3.9)), similar to what has been done in [9]–[12].

For $v = 1, 2, \dots, \sigma$ we put

$$\varphi_v(x) = x^\nu \exp(-\pi x^2), \quad \tilde{\varphi}_v(\xi) = i^{-\nu} \varphi_v(\xi), \quad \nu = \nu_v \in F_2. \quad (4.1)$$

Then

$$\Phi_v(\alpha; \omega_\nu) = G_\infty(\alpha + i\alpha_v + \nu), \quad \tilde{\Phi}_v(\alpha; \bar{\omega}_\nu) = i^{-\nu} G_\infty(\alpha - i\alpha_v + \nu), \quad (4.2)$$

where [13]

$$G_\infty(\alpha) = \int_{-\infty}^{\infty} |x|^{\alpha-1} \exp(-\pi x^2) dx = \pi^{-\alpha/2} \Gamma(\alpha/2).$$

Note owing to (2.10) that

$$\Gamma_\infty(\alpha; \nu) = i^{-\nu} \frac{G_\infty(\alpha + \nu)}{G_\infty(1 - \alpha + \nu)}, \quad \nu \in F_2. \quad (4.3)$$

For $v = \sigma + 1, \dots, \sigma + \tau$ we put

$$\varphi_v(x) = x^\nu \exp(-2\pi x\bar{x}), \nu \geq 0, \quad \varphi_v(x) = \bar{x}^{|\nu|} \exp(-2\pi x\bar{x}), \nu < 0,$$

$$\tilde{\varphi}_v(\xi) = i^{-\nu} \bar{\varphi}_v(\xi), \quad \nu = \nu_v \in Z.$$

Then

$$\begin{aligned} \Phi_v(\alpha; \omega_\nu) &= G_{-\infty}(\alpha + i\alpha_v + |\nu|/2), \\ \tilde{\Phi}_v(\alpha; \bar{\omega}_\nu) &= i^{-\nu} G_{-\infty}(\alpha - i\alpha_v + |\nu|/2) \end{aligned} \quad (4.4)$$

where [13]

$$G_{-\infty}(\alpha) = \int_{\mathbb{C}} (x\bar{x})^{\alpha-1} \exp(-2\pi x\bar{x}) dx = (2\pi)^{1-\alpha} \Gamma(\alpha).$$

Note owing to (2.15) that

$$\Gamma_{-\infty}(\alpha; \nu) = i^{-|\nu|} \frac{G_{-\infty}(\alpha + |\nu|/2)}{G_{-\infty}(1 - \alpha + |\nu|/2)}, \quad \nu \in Z. \quad (4.5)$$

For $v \in F_V$ we take function φ_v in form (2.22) (for field $\mathbb{K}_v$). Then owing to (2.20) we have

$$\Phi_v(\alpha; \omega_v) = \Gamma_v(\alpha + i\alpha_v; 1) = q_v^{(\alpha + i\alpha_v - 1/2)r_v} \Gamma_{q_v}(\alpha + i\alpha_v) \quad (4.6)$$

and owing to (2.8) and (4.6)

$$\tilde{\Phi}_v(\alpha; \bar{\omega}_v) = 1. \quad (4.7)$$

For $v \in R$ we take function φ_v in form (2.26) (for field $\mathbb{K}_v$). Then owing to (2.24) we have

$$\Phi_v(\alpha; \omega_v) = \Gamma_v(\alpha + i\alpha_v; \theta_v) = \kappa_v q_v^{(\alpha + i\alpha_v - 1/2)(r_v + \rho_v)}, \quad (4.8)$$

where a number $\kappa_v = \kappa(\theta_v)$ is defined by Eq. (2.25),

$$\kappa_v(\theta_v) = q_v^{\rho_v/2} \int_{|x|_v=1} \theta_v(x) \chi(\pi_v^{-r_v - \rho_v} x) d_v x, \quad |\kappa_v| = 1. \quad (4.9)$$

At last, owing to (2.8) and (4.8) we have

$$\tilde{\Phi}(\alpha; \bar{\omega}_v) = \theta_v(-1). \quad (4.10)$$

Substituting expressions (4.2), (4.4), (4.6)-(4.8) and (4.10) in the Tate formula (3.12) we get for all $V > \sigma + \tau$ the following Eq.

$$\begin{aligned} & \prod_{v=1}^{\sigma} G_{\infty}(\alpha + i\alpha_v + \nu_v) \prod_{v=\sigma+1}^{\sigma+\tau} G_{-\infty}(\alpha + i\alpha_v + |\nu_v|/2) \prod_{v \in F_V} q_v^{(\alpha + i\alpha_v - 1/2)r_v} \Gamma_{q_v}(\alpha + i\alpha_v) \\ & \times \prod_{v \in R} \kappa(\theta_v) q_v^{(\alpha + i\alpha_v - 1/2)(r_v + \rho_v)} L_V(\alpha; \omega) = \prod_{v=1}^{\sigma} i^{-\nu_v} G_{\infty}(1 - \alpha - i\alpha_v + \nu_v) \end{aligned}$$

$$\times \prod_{v=\sigma+1}^{\sigma+\tau} i^{-\nu_v} G_{-\infty}(1 - \alpha - \alpha_v + |\nu_v|/2) \prod_{v \in R} \theta_v(-1) L_V(1 - \alpha; \bar{\omega}). \quad (4.11)$$

Now we divide both sides of Eq. (4.11) on

$$\prod_{v=1}^{\sigma} G_{\infty}(1 - \alpha - i\alpha_v + \nu_v) \prod_{v=\sigma+1}^{\sigma+\tau} G_{-\infty}(1 - \alpha - i\alpha_v + |\nu_v|/2).$$

As a result, using Eqs (4.3),(4.5) and (3.7) (cf. [13],[10]), we get the following key formula for gamma-functions

$$\begin{aligned} \prod_{v=1}^{\sigma} \Gamma_{\infty}(\alpha + i\alpha_v; \nu_v) \prod_{v=\sigma+1}^{\sigma+\tau} \Gamma_{-\infty}(\alpha + i\alpha_v; \nu_v) \prod_{v \in F_V} \Gamma_{q_v}(\alpha + i\alpha_v) L_V(\alpha; \omega) \\ = \kappa \omega(C) [|D|N(J)]^{1/2-\alpha} L_V(1 - \alpha; \bar{\omega}) \end{aligned} \quad (4.12)$$

for all $V > \sigma + \tau$. In (4.12)

$$\kappa = \prod_{v=1}^{\infty} \kappa_v, \quad |\kappa| = 1, \quad (4.13)$$

where $\kappa_v = i^{-\nu_v}$, $v = 1, 2, \dots, \sigma$, $\kappa_v = i^{-|\nu_v|}$, $v = \sigma + 1, \dots, \sigma + \tau$, $\kappa_v = 1$, $v \in F$, $\kappa_v = \theta_v(-1) \bar{\kappa}(\theta_v)$, $v \in R$ (see (4.9)); and also

$$C = \prod_{v > \sigma + \tau} \pi_v^{r_v + \rho_v}, \quad \omega(C) = \prod_{v > \sigma + \tau} q_v^{-i\alpha_v(r_v + \rho_v)}. \quad (4.14)$$

As Euler's function $\Gamma(\alpha)$ does not vanish everywhere, so both sides of Eq. (4.12) are entire functions provided the set R is not empty.

Formula (4.12) is valid also for $V = \sigma + \tau + 1$ ($F_V = F$). In this case owing to (3.14) it gives the functional Eq. for the Dirichlet L -function [13],

$$\begin{aligned} \prod_{v=1}^{\sigma} \Gamma_{\infty}(\alpha + i\alpha_v; \nu_v) \prod_{v=\sigma+1}^{\sigma+\tau} \Gamma_{-\infty}(\alpha + i\alpha_v; \nu_v) L(\alpha; \omega) \\ = \kappa \omega(C) [|D|N(J)]^{1/2-\alpha} L(1 - \alpha; \bar{\omega}). \end{aligned} \quad (4.15)$$

By virtue of (3.13) for $\text{Re } \alpha < 0$

$$\lim_{V \rightarrow \infty} L_V(1 - \alpha; \bar{\omega}) = 1, \quad (4.16)$$

and thus by (4.12) there exists

$$\lim_{V \rightarrow \infty} \prod_{v \in F_V} \Gamma_{q_v}(\alpha + i\alpha_v) L_V(\alpha; \omega)$$

which we will denote by

$$\text{reg} \prod_{v \in F} \Gamma_{q_v}(\alpha + i\alpha_v), \quad (4.17)$$

, and this limit satisfies Eq.

$$\begin{aligned} & \prod_{v=1}^{\sigma} \Gamma_{\infty}(\alpha + i\alpha_v; \nu_v) \prod_{v=\sigma+1}^{\sigma+\tau} \Gamma_{-\infty}(\alpha + i\alpha_v; \nu_v) \text{reg} \prod_{v \in F} \Gamma_{q_v}(\alpha + i\alpha_v) \\ &= \kappa \omega(C) [|D| N(J)]^{1/2-\alpha}. \end{aligned} \quad (4.18)$$

For $\text{Re } \alpha \geq 0$ the regularization expression (4.17) is defined by Eq. (4.18) as the analytic continuation on α from domain $\text{Re } \alpha < 0$ (as a meromorphic function).

Eq. (4.18) is regularized adelic formula for gamma-functions of all completions of field $\mathbb{K}$.

For unramified characters ($\nu_v = 0, \alpha_v = 0, \kappa = 1, \omega(C) = 1, N(J) = 1$) Eq. (4.18) takes the form [9],[10]

$$\Gamma_{\infty}^{\sigma}(\alpha) \Gamma_{-\infty}^{\tau}(\alpha) \text{reg} \prod_{p=2}^{\infty} \prod_{j=1}^{m_p} \Gamma_{q_{pj}}(\alpha) = |D|^{1/2-\alpha} \quad (4.19)$$

where m_p is number of prime divisors of field $\mathbb{K}$ entering in decomposition of prime number p .

For field $\mathbb{K} = \mathbb{Q}$ ($\sigma = 1, \tau = 0, \alpha_\infty = 0, r_v = 0, D = 1$) Eq. (4.18) takes the form

$$\Gamma_\infty(\alpha; \nu) \operatorname{reg} \prod_{p \in F} \Gamma_p(\alpha + i\alpha_p) = \kappa \omega(C) N^{1/2-\alpha}(J), \quad (4.20)$$

where numbers $\nu, \alpha_p, \kappa, \omega(C), N(J)$ are defined by Eqs (3.6), (3.7), (4.13), (4.14). Note that owing to (3.6) and (1.5) in this case

$$\Gamma_p(\alpha + i\alpha_p) = \frac{1 - \theta(p)p^{\alpha-1}}{1 - \bar{\theta}(p)p^{-\alpha}}, \quad p = 2, 3, 5, \dots \quad (4.21)$$

§5. Regularized adelic formulas for beta-functions

Let characters ω and ω' be defined on $A^\times / \mathbb{K}^\times$. Suppose that ranks ρ_v, ρ'_v and ρ''_v of characters ω_v, ω'_v and $\omega''_v = \omega_v \omega'_v$ (of form (3.1)) of fields $\mathbb{K}_v$ are equal, $\rho_v = \rho'_v = \rho''_v$, for all $v \in R$. In this case the quantities $N(J)$ and C and the sets F and R turn out to be the same for the characters ω, ω' and ω'' .

Our goal is to prove the regularized formula (1.3) for beta-functions.

If we multiply Eq. (4.18) on the same Eq. in which magnitudes $\alpha, \alpha_v, \nu_v, \kappa_v$ and ω are changed by $\beta, \beta_v, \mu_v, \kappa'_v$ and ω' resp., and the Eq. obtained we divide on Eq. (4.18) in which just mentioned magnitudes are changed by $\alpha + \beta, \alpha_v + \beta_v, \nu_v + \mu_v, \kappa''_v$ and ω'' resp. As a result by using formulas (2.10), (2.12), (2.15), (2.17), (2.23) and (1.5) we get for $V > \sigma + \tau$ the following Eq.

$$\begin{aligned} & \prod_{v=1}^{\sigma} B_\infty(\alpha + i\alpha_v, \nu_v; \beta + i\beta_v, \mu_v) \prod_{v=\sigma+1}^{\sigma+\tau} B_{-\infty}(\alpha + i\alpha_v, \nu_v; \beta + i\beta_v, \mu_v) \\ & \times \prod_{v \in F_V} B_{q_v}(\alpha + i\alpha_v, \beta + i\beta_v) L_V(\alpha; \omega) L_V(\beta; \omega') L_V^{-1}(\alpha + \beta; \omega'') \\ & = \kappa \sqrt{|D|N(J)} L_V(1 - \alpha; \bar{\omega}) L_V(1 - \beta; \bar{\omega}') L_V^{-1}(1 - \alpha - \beta; \bar{\omega}''), \quad (5.1) \end{aligned}$$

where

$$\kappa = \prod_{v \in R} \kappa_v \kappa'_v \bar{\kappa}_v'', \quad |\kappa| = 1. \quad (5.2)$$

Passing on to limit as $V \rightarrow \infty$ in Eq. (5.1) for $\operatorname{Re} \alpha < 0$ and $\operatorname{Re} \beta < 0$, using (4.16) and denoting

$$\begin{aligned} \lim_{V \rightarrow \infty} \prod_{v \in F_V} B_{q_v}(\alpha + i\alpha_v, \beta + i\beta_v) L_V(\alpha; \omega) L_V(\beta; \omega') L_V^{-1}(\alpha + \beta; \omega'') \\ = \operatorname{reg} \prod_{v \in F} B_{q_v}(\alpha + i\alpha_v, \beta + i\beta_v), \end{aligned} \quad (5.3)$$

we get Eq.

$$\begin{aligned} \prod_{v=1}^{\sigma} B_{\infty}(\alpha + i\alpha_v, \nu_v; \beta + i\beta_v, \mu_v) \prod_{v=\sigma+1}^{\sigma+\tau} B_{-\infty}(\alpha + i\alpha_v, \nu_v; \beta + i\beta_v, \mu_v) \\ \times \operatorname{reg} \prod_{v \in F} B_{q_v}(\alpha + i\alpha_v, \beta + i\beta_v) = \kappa \sqrt{|D|N(J)}. \end{aligned} \quad (5.4)$$

So we have just obtained the following

Theorem. *Let $\mathbb{K}$ be an algebraic numbers field of degree $n = \sigma + 2\tau$, ω, ω' and $\omega'' = \omega\omega'$ be characters on $A^\times / \mathbb{K}^\times$ and their local ranks ρ_v, ρ'_v and ρ''_v be equal, $\rho_v = \rho'_v = \rho''_v, v \in R$. Then for beta-functions of all completions of field $\mathbb{K}$ the regularized adelic formula (5.4) is valid.*

For unramified characters formula (5.4) takes the form [1],[2]

$$B_{\infty}^{\sigma}(\alpha, \beta) B_{-\infty}^{\tau}(\alpha, \beta) \operatorname{reg} \prod_{p=2}^{\infty} \prod_{j=1}^{m_p} B_{q_{pj}}(\alpha, \beta) = \sqrt{|D|}. \quad (5.5)$$

For field $\mathbb{K} = \mathbb{Q}$ formula (5.4) takes the following explicit form

$$B_{\infty}(\alpha, \nu; \beta, \mu) \operatorname{reg} \prod_{p \in F} B_p(\alpha + i\alpha_p, \beta + i\beta_p) = \kappa \sqrt{N(J)} \quad (5.6)$$

where owing to (3.6) and (3.5)

$$\theta(-1) = \theta'(-1) = 1, \quad p^{i\alpha_p} = \theta(p), \quad p^{i\beta_p} = \theta'(p), p \in F. \quad (5.7)$$

Similar, in a more complicated way, regularized adelic formulas can be written out for cases when local ranks $\rho_v, \rho'_v, \rho''_v$ of characters $\omega, \omega', \omega''$ are different.

References

- 1 P. G. O. Freund and E. Witten, Adelic String Amplitudes, Phys. Lett. B, 199:191–194 (1987).
- 2 I. V. Volovich, Harmonic Analysis and p-Adic Strings, Lett. Math. Phys., 16:61–67 (1988).
- 3 P. G. O. Freund and M. Olson, Non-Archimedean Strings, Phys. Lett. B, 199:186–190 (1987).
- 4 I. V. Volovich, p-Adic String, Class. Quantum Grav., 4:L83–L87 (1987).
- 5 I. Ya. Aref'eva, B. G. Dragović and I. V. Volovich, On the Adelic String Amplitudes, Phys. Lett. B, 209:445–450 (1988).
- 6 I. Ya. Aref'eva, B. G. Dragović and I. V. Volovich, Open and Closed p-Adic Strings and Quadratic Extensions of Number Fields, Phys. Lett. B, 212:283–291 (1988).
- 7 P. H. Frampton, Y. Okada and M. R. Ubriaco, On Adelic Formulas for p-Adic String, Phys. Lett. B, 213:260–262 (1988).
- 8 P. H. Frampton, and H. Nishino, Theory of p-Adic Closed Strings, Phys. Rev. Lett., 62:1960–1963 (1989).
- 9 V. S. Vladimirov and T. M. Sapuzhak, Adelic Formulas for String Amplitudes in Fields of Algebraic Numbers, Lett. Math. Phys., 37(2):233–242 (1996).
- 10 V. S. Vladimirov, Adelic Formulas for Gamma- and Beta-functions of Completions of Algebraic Number Fields and Their Applications to String Amplitudes, Izv. RAN, ser. math., 60, № 1:63–86 (1996)(in Russian).

11. V. S. Vladimirov, On the Freund-Witten Adelic Formula for Veneziano Amplitudes, Lett. Math. Phys., 27:123–131 (1993).
12. V. S. Vladimirov, The Adelic Freund-Witten Formulas for the Veneziano and Virasoro-Shapiro Amplitudes, Uspechi Math. Nauk., 48, № 6:3–38 (1993) (in Russian).
13. A. Weil. Basic Numbers Theory, Springer, Berlin etc, 1967.
14. Z. I. Borevich and I. R. Shafarevich. Théory de nombres, Gauthier-Villars, Paris, 1967.
15. S. Lang. Algebraic Numbers, Adisson-Wesley, London, 1964.
16. W. H. Schikhof. Ultrametric Calculus. An Introduction to p-Adic Analysis, Cambridge Univ. Press, 1984, XI+306p.
17. N. Koblitz. p-Adic Numbers, p-Adic Analysis, and Zeta-functions, Springer, Berlin etc, 1977.
18. I. M. Gel'fand, M. I. Graev and I. I. Pjatetskii-Shapiro. Representations Theory and Automorphic Functions, Saunders, Philadelphia, 1969.
19. V. S. Vladimirov, I. V. Volovich and E. I. Zelenov. p-Adic Analysis and Mathematical Physics, World Scientific, Singapore, 1994, XVIII+319p.
20. Andrei Khrennikov. p-Adic Valued Distributions in Mathematical Physics, Kluver, Dordrecht etc, 1994, XIV+264p.
21. V. S. Vladimirov. Adelic Formulas for Gamma- and Beta-functions in Fields of Algebraic Numbers, Doklady Mathematics, 347(1):11–15 (1996).